

DECOMPRESSION ILLNESS (Dysbarism)

APPROVED: Gregory Gilbert, MD EMS Medical Director
 Nancy Lapolla EMS Director

DATE: July 2018

Information Needed:

- Length of time patient submerged (total dive time)
- Number of dives made
- Duration of and time since descent/ascent (total surface interval)
- Were decompression dives made and at what interval?
- Depth of deepest submersion and depth of last dive
- Any loss of consciousness
- Temperature of the water
- Mechanism of injury suggestive of head/neck injury
- Did the diver perform an emergency ascent? If so, from what depth?
- Was the dive made with compressed air or mixed gases?
- Was there any air flight in last 24 hours?

Objective Findings:

- Joint pain (location)
- Pulmonary exam: rales or signs of pulmonary edema, respiratory distress
- Neurologic exam: monitor continuously for evolution of focal deficits
- Cardiac rhythm
- Resolution of symptoms with O₂

Treatment:

- Routine medical care
- 100% oxygen by non-rebreather mask
- Place patient in left lateral position, have suction ready
- Immediate transport to appropriate local hospital
- IV access, treat for dehydration

For joint pain:

- Consider pain control, see Interim Adult Interim Pain Assessment and Management protocol (June 2018).

Precautions and Comments:

- Shock position is contraindicated (this position does not affect circulatory system, emboli, and it may increase cerebral edema)
- Keep patient supine/left lateral recumbent position if possible to reduce risk of cerebral air embolism

- Focal deficits and neurologic changes indicate need for rapid transport to a hyperbaric chamber after stabilization at a local emergency department
- Low altitude, rapid air medical evacuation if necessary to the nearest decompression chamber
- Rapid ascent or breath-holding during ascent may cause central air embolism or ruptured tympanic membrane(s)
- Be alert for recurring hypoxia
- O₂ may resolve symptoms but is not a substitute for definitive care
- As much as 50% of dive injuries occur at depths not needing recompression
- Dysbarism may also occur in workers conducting operations in a pressurized air environment, e.g. bridge caisson workers. Consult on-scene work supervisors
- DAN (Divers Alert Network) Diving Emergency Hotline 919-684-9111 (24 hours/day)