

## OBSTETRICAL/GYNECOLOGICAL EMERGENCIES/CHILDBIRTH

APPROVED: Gregory Gilbert, MD EMS Medical Director  
Nancy Lapolla EMS Director

DATE: July 2018

### **Vaginal Bleeding**

#### **Subjective Findings:**

- Abdominal Pain
- Estimate Blood Loss
  - Duration of bleeding
  - Quality of bleeding
  - Number of pads per hour
- Last menstrual period
- For known pregnancy – How many weeks/months?
- Estimated due date

#### **Objective Findings:**

- Blood Pressure
- Vaginal bleeding
  - with possible products of conception: Collect and transport with the patient
- Observe for signs of shock

#### **Treatment:**

- Consider second IV access
- If hypotensive (SBP < 90 or signs of poor perfusion), fluid challenge of 250-1000 ml NS IV. If SBP remains < 90 continue fluid resuscitation. Titrate to SBP of 90 or symptoms of improved perfusion
- Keep NPO
- Place pad or large dressing over vaginal opening if bleeding
- Save and transport all tissue or fetal remains
- If pregnant and over 20 weeks, transport to facility with OB capabilities and don't encourage delivery

### **Pre-eclampsia or Eclampsia**

#### **Objective Findings:**

- Hypertension (BP > 140/90, or SBP > 30 mm Hg above baseline, or DBP > 15 mm Hg above baseline)
- Over 20 weeks pregnant
- Altered mental status, blurred vision, headache, or seizures
- Pedal edema

#### **Treatment:**

- High flow oxygen
- IV access
- Minimize stimulation (lights, noise, other stressors)

- Left lateral decubitus position
- For seizures, midazolam (Versed®):
  - 1-2 mg IV/IO. May repeat every 5 minutes, up to a maximum dose of 10 mg
  - 1-5 mg IN. May repeat in 10 minutes up to a maximum dose of 10 mg.
  - Monitor the patient's EKG monitor and pulse oximetry after administration.

## **Childbirth**

### **Information Needed:**

- Estimated due date, last menstrual period
- Anticipated problems (multiple fetuses, premature delivery, placenta previa, lack of prenatal care, use of narcotics or stimulants, etc.)
- Gravida/para
- Onset of regular contractions
- Rupture of membranes, color of fluids, time of rupture
- Frequency and duration of contractions
- Urge to bear down or have bowel movement

### **Objective Findings:**

Observe perineal area for:

- Fluid or bleeding
- Crowning (check during contraction)
- Abnormal presentation (breech, extremity, cord)

#### **Treatment:**

- Routine medical care
- Oxygen as indicated
- If birth is not imminent, place patient in left lateral position
- IV access when appropriate

## **Normal Delivery**

### **Treatment:**

- Assist with delivery
- Clean, preferably sterile technique
- Control and guide delivery of baby's head and body
- Check for nuchal cord - slide overhead if possible: If tight, double clamp and cut, unwind, then deliver baby quickly
- Keep the baby at or below the level of the mother until the cord is clamped
- Suction mouth, then nares
- Double clamp and cut cord
- Dry and wrap infant for warmth (especially the head); if possible place with the mother for shared body heat
- Note time of delivery

- Assess infant's status, respirations and pulse rate (APGAR if possible)
- Evaluate mother post-delivery for excessive bleeding

### **Post-partum hemorrhage**

#### Treatment:

- Fundal massage
- Fluid challenge, 250-1000 ml Normal Saline

### **Breech Delivery**

#### Treatment:

- Provide airway with gloved hand for baby if needed
- If unable to deliver or delivery is not imminent, place patient in left lateral Trendelenburg position and transport rapidly
- In the setting of a limb presentation, do not pull on the delivered extremity

### **Prolapsed Cord**

#### Treatment:

- Left lateral Trendelenburg position and elevate hips when possible
- If the cord is pulseless, manually displace presenting part above the cord
- Avoid manipulation of the cord, cover with a moist sterile dressing, and try to keep cord warm by keeping it close to the patient's body
- Rapid transport

### **Precautions and Comments:**

- Ectopic pregnancy should be considered in all women of childbearing age with either abdominal pain or vaginal bleeding.
- Eclampsia can occur up to a week after delivery
- The first priority in childbirth is assisting the mother with delivery of the child
- The primary enemy of a newborn is hypothermia and can occur within minutes
- Consider early suctioning after delivery of the infant with evidence of meconium
- Do not pull on the cord
- Do not delay transport for delivery of placenta. If placenta delivers, then place it in a plastic bag for transport with the mother
- If hemorrhage is present prior to delivery in a patient who is near term, consider emergencies such as placental abruption, placenta previa, and uterine rupture. Treat with IV fluids (consider a second IV) and rapid transport to an appropriate hospital
- Massage fundus after delivery of placenta